

Malignant T cells express lymphotoxin α and drive endothelial activation in cutaneous T cell lymphoma

Supplementary Material




Figure S1: LT α is not expressed in benign inflammatory skin disorders. Representative frozen biopsy from a patient with benign skin inflammation was subjected to IHC with an antibody directed against LT α showing negative staining of perivascular small lymphocytes. Essentially similar results were found in 9 out of 9 cases.




Figure S2: JAK regulate $LT\alpha$ expression in malignant CTCL T cells. Two malignant CTCL T cell lines (Mac-2a and PB2B) were incubated with 10μM CP6990,550 for 24h. Subsequent the concentrations of $LT\alpha$ in the culture supernatants were measured by an $LT\alpha$ specific ELISA. Bars represent mean values of three independent experiments. * $p < 0.05$ compared to control (paired t-test).




Figure S3: STAT3 do not regulate the LTα expression in CTCL. MyLa2059 cells were transiently transfected with STAT3 siRNA. 24 hours post-transfection, supernatants were harvested and LTα concentration measured by ELISA. Bars represent mean values of three independent experiments.




Figure S4: STAT5 bind to the LT α promoter region. Pull-down assay using oligonucleotides representing a STAT binding site in the promoter region of the LT α gene. STAT5 bind to the sequence, while STAT3 do not.




Figure S5: *LT α* regulates the IL-6 expression in malignant CTCL T cells. The IL-6 expression in two malignant CTCL cell lines (Mac-2a and PB2B) was measured by ELISA following incubation with 100ug/ml Etanercept for 24 h. Bars represent one experiment.